

EFFECTS OF SECONDARY STATIC STRESS TRIGGERING ON THE SPATIAL DISTRIBUTION OF AFTERSHOCKS, A CASE STUDY, 2003 BAM EARTHQUAKE (SE IRAN)

Behnam MALEKI ASAYESH

Ph.D. Candidate, IIEES, Tehran, Iran
b.malekiasayesh@iiees.ac.ir

Hamid ZAFARANI

Associate Professor, IIEES, Tehran, Iran
h.zafarani@iiees.ac.ir

Mohammad TATAR

Associate Professor, IIEES, Tehran, Iran
mtatar@iiees.ac.ir

Keywords: Coulomb stress, Static triggering, Aftershocks, Nodal plane, Bam Earthquake, Iran

Immediate after a large earthquake, accurate prediction of spatial and temporal distribution of aftershocks has a great importance for planning search and rescue activities. Currently, the most sophisticated approach to this goal is probabilistic aftershock hazard assessment (PASHA). Spatial distribution of the aftershocks following moderate to large earthquakes correlate well with the imparted stress due to the mainshock. Furthermore the secondary static stress changes caused by smaller events (aftershocks) could have effect on the triggering of aftershocks and should be considered in the calculations. The 26 December 2003 (Mw 6.6) Bam earthquake with more than 26000 casualties is one of the most destructive events in the recorded history of Iran. This earthquake was an interesting event and was investigated in a majority of aspects.

One of the successful models that could forecast the spatial distribution of the aftershocks and next mainshocks is Coulomb stress changes. Aftershocks and subsequent mainshocks often occur in regions that experienced an increase in Coulomb stress caused by the mainshock, and earthquakes become less prevalent than before the main shock in regions subjected to a Coulomb stress drop. Good variable-slip fault model and precise aftershocks data enabled us to impart Coulomb stress changes due to mainshock and secondary static stress triggering on the nodal planes of aftershocks to learn whether they were brought closer to failure.

We used recently published high-quality focal mechanisms and hypocenters to reassess the role of small to moderate earthquakes for static stress triggering of aftershocks during the Bam earthquake. By imparting Coulomb stress changes due to the mainshock on the nodal planes of the 158 aftershocks we showed that 77.8% (123 from 158) of the aftershocks received positive stress changes at least in one nodal plane (Figure 1). We also calculated Coulomb stress changes imparted by the mainshock and aftershocks ($1 \leq M \leq 4.1$) onto subsequent aftershocks nodal planes and found that 81.6% (129 of 158) of aftershocks received positive stress changes at least in one nodal plane. In summary, 77.8% of aftershocks are encouraged by the main shocks, while adding secondary stress encourages 81.6%. Therefore, by adding secondary stress the Coulomb Index (CI), the fraction of events that received net positive Coulomb stress changes compared to the total number of events, increased from 0.778 to 0.816.

REFERENCES

- Asayesh, B.M., Hamzehloo, H., and Zafarani, H. (2018). Coulomb stress changes due to main earthquakes in Southeast Iran during 1981 to 2011. *Journal of Seismology*, 1-16.
- Funning, G.J., Parsons, B., Wright, T.J., Jackson, J.A., and Fielding, E.J. (2005). Surface displacements and source parameters of the 2003 Bam (Iran) earthquake from Envisat advanced synthetic aperture radar imagery. *Journal of Geophysical Research: Solid Earth*, 110(B9).



Jackson, J., Bouchon, M., Fielding, E., Funning, G., Ghorashi, M., Hatzfeld, D., Nazari, H., Parsons, B., Priestley, K., Talebian, M., Tatar, M., Walker, R., and Wright, T. (2006). Seismotectonic, rupture process, and earthquake-hazard aspects of the 2003 December 26 Bam, Iran, earthquake. *Geophysical Journal International*, 166(3), 1270-1292.

King, G.C., Stein, R.S., and Lin, J. (1994). Static stress changes and the triggering of earthquakes. *Bulletin of the Seismological Society of America*, 84(3), 935-953.

Ma, K.F., Chan, C.H., and Stein, R.S. (2005). Response of seismicity to Coulomb stress triggers and shadows of the 1999 Mw= 7.6 Chi-Chi, Taiwan, earthquake. *Journal of Geophysical Research: Solid Earth*, 110(B5).

Tatar, M., Hatzfeld, D., Moradi, A.S., and Paul, A. (2005). The 2003 December 26 Bam earthquake (Iran), Mw 6.6, aftershock sequence. *Geophysical Journal International*, 163(1), 90-105.

Toda, S., Stein, R.S., Reasenber, P.A., Dieterich, J.H., and Yoshida, A. (1998). Stress transferred by the 1995 Mw= 6.9 Kobe, Japan, shock: Effect on aftershocks and future earthquake probabilities. *Journal of Geophysical Research: Solid Earth*, 103(B10), 24543-24565.




Figure 1. Coulomb stress changes on the nodal plane of the Bam aftershocks. a) Imparted stress on the 81 aftershocks that are put in category A. b) Imparted stress on the 68 and 9 aftershocks that are put in category B and C, respectively.